



BACKYARDS
for *Butterflies*

OHIO DEPARTMENT OF NATURAL RESOURCES DIVISION OF WILDLIFE
BACKYARDS FOR BUTTERFLIES

TABLE OF CONTENTS

LIFE CYCLE.....	03	SUMMER BACKYARDS..	08
GARDENING TIPS	04	FALL BACKYARDS	10
GLOSSARY.....	04	WINTER BACKYARDS....	12
GET STARTED	05	NECTAR PLANTS.....	14
SPRING BACKYARDS	06	BUTTERFLIES.....	16

FEATURED BUTTERFLIES


TIGER SWALLOWTAIL	16	AMERICAN LADY	23
SPICEBUSH SWALLOWTAIL.....	17	MONARCH	24
GIANT SWALLOWTAIL	18	GREAT SPANGLED FRITILLARY	25
BLACK SWALLOWTAIL	19	NORTHERN PEARLY EYE	26
PEARL CRESCENT	20	SILVER SPOTTED SKIPPER	26
RED ADMIRAL	21	SPRING AZURE	27
QUESTION MARK	22	MILBERT'S TORTOISESHELL ...	27


INTRODUCTION

More than 140 different kinds of butterflies have been recorded in Ohio. You won't find all of them in your yard, but more than two dozen species, ranging in size from the tiny blue spring azure to the magnificent monarch, can be found. If you want to attract wildlife to your property, start with butterflies! As more wildlife species that were once common are now in decline there is a greater sense of urgency for Ohioans to take action. Population declines have been seen across all different types of animals including insects, reptiles, amphibians, birds and mammals. The sheer amount of habitat loss needs to be reversed. One of the best opportunities is to provide habitat in your own yard. Providing habitat for wildlife means providing water, shelter and food. It is easy to do with a few simple steps, so let's get started today and begin your journey to help conserve Ohio's wildlife.

ACKNOWLEDGMENTS

Special thanks to Jeff Burriss, Rick Gardner, Jamey Emmert, Kate Parsons, Lindsay Rist, Erica Eppert, Jerry Wiedmann (Ohio Lepidopterists), Cheryl Harner (North Central Ohio Land Conservancy), and Denise Ellsworth (OSU) for their contributions.





BUTTERFLY LIFE CYCLE

The life cycle of butterflies may seem straight forward but many species have interesting variations on the theme called “complete metamorphosis.” Some butterfly species complete the cycle several times in one year, others only once. Some species overwinter as eggs, some as larvae, some as pupae in the chrysalis and some as adults. Most adult butterflies are short-lived. Few live more than 7-10 days, with the exception of overwintering species, such as monarchs, mourning cloaks, and question marks, which may live as long as 12 months.

WHAT YOU CAN DO

- 🦋 Provide habitat for wildlife year-round. For example, plan your garden so you have flowering plants from early spring to late fall. Consider what butterflies need to complete their entire life cycle – nectar plants, host plants, and areas for overwintering.
- 🦋 Plant native plants whenever possible. Native plants provide butterflies and their caterpillars the resources they need and when they need them. For example, don’t plant tropical milkweed (*Asclepias curassavica*) which is not native to Ohio when there are many native milkweed species available.
- 🦋 Many perennial plants won’t bloom the first year. Have patience for blooms.
- 🦋 Need to start small? No effort is too small to help butterflies. Even container gardens can help provide blooms they need for nectar or host plants (food) for caterpillars.
- 🦋 Attracting pollinators to your yard will help ensure pollination for your vegetable garden.
- 🦋 Leaving the leaves on the ground and dead stems through the winter is important. Caterpillars and butterflies use these for shelter to survive the freezing temperatures
- Remove invasive plants which disrupt natural cycles and compete with native plants for resources.
- 🦋 Pesticides, an umbrella term for insecticides, herbicides and fungicides, can be in your seeds, plants or soil that you purchase. Even products labeled organic can be toxic to butterflies. Ask before you buy.
- 🦋 Make special enhancements for the butterflies you want to attract – water, feeders with rotting fruit, or wet sandy areas for puddling.
- 🦋 Expect insect damage – native plants can typically tolerate use by insects.
- 🦋 If you have a large site that you are restoring, contact the Division of Wildlife Private Lands Program at 1-800-WILDLIFE (945-3543) for guidance and resources.

BASIC GARDENING TIPS

Planting a wide range of nectar and host plants is the best strategy for attracting a wide assortment of butterfly species. A large patch of bright colorful flowers might be appealing to butterflies, but they will linger if there are also areas that meet additional essential needs for shelter, water, and sun.

Create vital plant diversity by choosing plants of different types. Think about shrubs, trees, perennials, and even vines. Select plants that grow to different heights, with a variety of flower shapes and colors that bloom at different times (bloom periods) so there are always offerings of nectar. If the garden is small with little space for trees or shrubs, consider an arbor covered with vines to create height and interest.

Aim for basic gardening success by reading the tag that comes with the plant or ask your local nursery expert for guidance. Following plant preferences will help achieve the best results!

----- SUN -----

Pick the right spot. Many flowers that are attractive to butterflies and the butterflies themselves often prefer access to several hours of direct sun.

----- SOIL -----

Be mindful of your garden location and the soil type. Do some digging at home and check out a shovel-full of soil before shopping for plants. Some species will thrive in wet areas while others thrive in the driest of soil. Choose plants according to the characteristics of the space in which they will be growing.

----- SIZE -----

Carefully select the flower species according to growth. Plant flowers likely to grow the tallest in the middle or the back of the garden and the shorter, ground-covers around the edges.

----- SPACE -----

Try not to pack the garden too densely with flowers. Be patient. It won't be long before plants will spread and start filling in the gaps.

GLOSSARY

ANNUAL - Plants with a one-year life cycle. They grow, bloom, produce seeds, and die in one year's time.

BROOD - A generation of butterflies that completes a whole life cycle.

BUTTERFLY - a nectar-feeding, diurnal insect with two pairs of large, typically brightly colored wings that are covered with microscopic scales. Distinguished from moths by having clubbed antennae, holding their wings erect when at rest.

CATERPILLAR - the larva of a butterfly or moth, having a segmented wormlike body with three pairs of true legs and several pairs of appendages similar to legs. Caterpillars may be hairy, have warning coloration, or be colored to resemble their surroundings.

CHRYsalis - a hardened coating of protein protecting an insect pupa inside; mostly associated with butterflies

COCOON - a silk casing, protecting an insect pupa inside; mostly associated with moths

DIAPAUSE - a period of suspended development, especially during unfavorable environmental conditions

DIURNAL - of, relating to, or occurring during the day

FORB - a herbaceous flowering plant other than a grass

FUNGICIDE - a substance that destroys fungus (molds, yeast, mushrooms, and toadstools)

HERBICIDE - a substance that is toxic to plants, used to destroy unwanted vegetation

HOST PLANT - known as nurseries of the garden; specific plants on which caterpillars must eat

INSECTICIDE - a substance used for killing insects

LARVA - the active immature form of an insect

LOAM - a fertile soil with roughly equal proportions of sand, silt, and clay

NATIVE - plants indigenous to a given area in geologic time

NECTAR - a sugar-rich liquid produced by plants in glands called nectaries

NOCTURNAL - of, relating to, or occurring at night

PERENNIAL - Plants that persist for many growing seasons. Generally the top portion of the plant dies back each winter and regrows the following spring from the same root system

PESTICIDE - a substance used for destroying insects or other organisms

PRUNING - horticultural and silvicultural practice involving the selective removal of certain parts of a plant, such as branches, buds, or roots


PUPA - the intermediate stage between the larva and adult insect

SKIPPER - a diurnal insect related to butterflies. Instead of knobbed antennae, the tips are usually hooked and the wing structure differs from butterflies. Flight is more erratic than most butterflies. Typically brown, tan, or yellow-orange in color.

GET YOUR WILDLIFE HABITAT CERTIFIED

Anyone can create a welcoming haven for local wildlife. The National Wildlife Federation is offering habitat enthusiasts the opportunity to show your commitment to wildlife. Turn your yard, balcony container garden, schoolyard, work landscape, or roadside greenspace into a Certified Wildlife Habitat® and make a lasting difference for Ohio's wildlife.

nwf.org/garden-for-wildlife/certify



GET STARTED

This guide presents a seasonal approach to building, maintaining, and enjoying a productive butterfly garden. Like all of nature, butterfly activity is influenced by the seasons, so your plans for gardening should be as well. Whether seeking host plants for egg laying, nectaring on their favorite blossoms, or finding a place to hibernate, your garden can be a haven to caterpillars and butterflies throughout the year. In the spring, gently clean up the garden while being mindful of sleeping caterpillars and the few species that have overwintered as adults. This helps promote a flurry of butterfly action as the summer sun awakens their activity levels for feeding and reproducing. The peak slowly fades into fall when some species by now have perished and others are making their flights southward to warmer climates.



SPRING

PLANTING, SEEDING, & MAINTENANCE

Postpone spring cleaning - delayed mowing allows grasses and forbs to mature and flower, providing much needed pollen, nectar, and seed resources to pollinators. Additionally, mature plants provide habitat needed for nesting, egg laying, and shelter. In early spring, many butterflies are still in diapause (hibernation), therefore they're basically still sleeping. Lots of beneficial insects spend the winter hunkered down in hollow plant stems either as adults or pupae. Cutting down the dead plant stems too early in the spring will disturb them before they have a chance to emerge.

When you do clean up, toss cut stems onto the compost pile or spread them near the edge of the woods. Many of the insects taking shelter inside the plant stems will still be able to emerge when the time is right. Another option is to bundle cut stems and hang them on a fence or lean them against a tree. The insects sheltering inside of them will emerge when they're ready. Bees might also use the stems as brood chambers all summer long.

BACKYARD PROJECTS FOR SPRING

Prune with care. Some moths and butterflies spend the winter in a delicate cocoon dangling from a branch, including the swallowtails, sulfurs, and spring azures. Allow any branches with a cocoon or chrysalis present to stay intact. You can always cut them back later in the season.

FIND A NURSERY

The Lake Erie Allegheny Partnership for Biodiversity provides a nursery map tool to help gardeners find native plants for sale.

<http://leapbio.github.io/nurseries/>

SWALLOWTAILS ON PUCCOON
Photo by NINA HARFMAN

COMMON SPRING BLOOMS



CANADA ANEMONE

Anemone canadensis

In only a few seasons, a few plants can quickly become a beautiful mat of white flowers. Even when not in bloom, the foliage is quite attractive. Prefers damp soil of sand, loam, or clay with full sun or partial shade. Height 1-2 feet. This plant can form dense colonies and may crowd out smaller vegetation.



GOLDEN ALEXANDER

Zizia aurea

Golden Alexanders is a member of the carrot family (*Apiaceae*). Bright yellow flowers bring color in early spring and are very attractive to bees. This is an adaptable plant for average soils and moisture as well as full sun or partial shade. Height 2.5-3.5 feet.



WILD COLUMBINE

Aquilegia canadensis

Blooms in a variety of colors including yellow and red. The bell-shaped flowers attract hummingbirds. Columbines will readily multiply once established. Wild columbine grows best in partial shade but can succeed in full sun or shade and is tolerant of a wide range of soils. Height 1-3 feet.



WILD GERANIUM

Geranium maculatum

Blooms of pale pink, deep pink and lilac show from April through May. It attracts beneficial insects, songbirds, and butterflies. Easily grown in average or medium, well-drained soil in full sun to part shade. Prefers moist soils, but tolerates poor soils. Height 1.5-2 feet.

SUMMER

PLANTING, SEEDING, & MAINTENANCE

One of the best things about planting native is how hands-off you can be with your garden. Native plants are adapted to the local climate and soil conditions where they naturally occur so there's little need to water during the driest of conditions.

Don't be afraid to "train" plants to grow at the width and height you desire during the blooming season. Just be judicious so as not to remove too many buds/flowers at once.

Maintain an adequate supply of host plants. Caterpillars are voracious eaters. Monitor the supply of host plants in your yard and add plants as needed.

BACKYARD PROJECTS FOR SUMMER

Create mineral sources. A small patch of wet soil will attract butterflies seeking minerals. A shallow dish with rocks or pebbles covered halfway with water will create a butterfly puddling site. During the summer, this requires some vigilance because water evaporates quickly. Pollinators will visit the dish to take in trace minerals from the water. Untreated tap water works fine here, because chlorine dissipates into the air after 24 hours. A garden pond lined with flat rocks is a permanent alternative.

iNATURALIST

iNaturalist is a citizen science project and online social network of naturalists, citizen scientists, and biologists built on the concept of mapping and sharing observations of biodiversity across the globe. iNaturalist may be accessed via its website or from its mobile applications. Record your observations, share with fellow naturalists, and discuss your findings!

inaturalist.org

RED ADMIRAL ON BLAZING STAR
Photo by M. CHUPPICH

COMMON SUMMER BLOOMS



BEE BALM

Monarda didyma

Prefers rich, humusy soils in full sun, although soil should not be allowed to dry out. Great for attracting hummingbirds! Deadhead flowers to prolong blooms. Divide clumps every 3-4 years to prevent overcrowding and to control spread. Good air circulation will help combat fungal leaf diseases. Height 2-4 feet.



DENSE BLAZING STAR

Liatris spicata

Easily grown in average or medium, well-drained soils in full sun. Somewhat tolerant of poor soils, but prefers moist, fertile ones and generally performs better in moist soils than most other species of *Liatris*. Intolerant of wet soils in winter. Tolerant of summer heat and humidity. Height 2-4 feet.



COMMON MILKWEED

Asclepias syriaca

A native perennial with clusters of upturned deep pink flowers. The leaves are a critical food source for monarch caterpillars and are very resistant to deer. Flowers are followed by seed pods which open and release silky-haired seeds. Grows in moist to mesic soils in full sun or partial shade. Height 2.5-5 feet.



PURPLE CONEFLOWER

Echinacea purpurea

Easily grown in average, dry to medium, well-drained soil in full sun to part shade. Best in full sun. Tolerant of drought, heat, humidity and poor soil. Divide clumps every 4 years to prevent overcrowding. Plants usually rebloom without deadheading. Self-seeds if some of the seed heads are left in place. Height 2-4 feet.



COMMON BUCKEYES ON ASTER
Photo by JOSHUA RICE

FALL

PLANTING, SEEDING, & MAINTENANCE

Winter habitat comes in the form of stems, leaves, and debris from your plants. You should leave these in place and try to avoid cleaning up flower beds and borders in the fall. Many native bees and butterflies overwinter on or inside the stems, and birds take shelter from harsh winter winds in the cover this debris provides. Toads nestle down in leaf debris and under loose mulch.

You may not want to display twig or stone piles right in the center of your garden, but tucking them away behind shrubs or a shed, out of sight and undisturbed, makes for a perfect insect-friendly hideaway.

Collect milkweed seeds and other native seeds. Some species need the cold temperatures of the winter months in order to germinate correctly, you can recreate this artificially by cold stratifying them in your refrigerator.

BACKYARD PROJECTS FOR FALL

Overwintering butterflies need cover. This may already be present in the habitat or nearby. Because species may overwinter in any of the four stages, a variety of cover is needed. Butterflies overwintering in the adult stage may use the peeling bark of trees, perennial plants, old logs or fences and firewood piles. Similar sites are used by overwintering pupae. Butterflies overwintering as caterpillars or eggs use herbaceous perennials, shrubs, and trees (see the Spring project section)

COMMON FALL BLOOMS



WHITE WOOD ASTER

Eurybia divaricata

This aster of delicate, airy clouds of white daisy-like flowers forms colonies and blooms in late summer and early fall. The small, white flowers show a yellow center that fade to red atop dark green to black stems. White wood aster grows best in partial to full shade with mesic to dry soils. Height 1-3 feet.



BLUE VERVAIN

Verbena hastata

Plants thrive in full or part sun in moist or wet mucky soils. It blooms from midsummer to early autumn. This plant adapts readily to degraded wetlands and other disturbed areas, but it can be found in higher quality habitats as well. Height 2-6 feet.



LITTLE BLUE STEM

Schizachyrium scoparium

Little Blue Stem grass displays a blue-green color in the summer turning shades of brown, copper, and crimson that will remain all winter. It is very drought-tolerant, but can do well in mesic situations. The grass gets fluffy white seed heads. Highly deer resistant and can grow in full sun or partial shade. Height 2-4 feet.



NATIVE ASTERS

Symphotrichum spp.

Most asters are late-blooming perennials that provide nectar to migrating species, like monarchs and painted ladies. For wintering species, aster nectar can help build up the stores of carbohydrates needed to survive their hibernation. Full sun and well-drained soils are ideal. Height 3-6 feet.



WINTER

PLANTING, SEEDING, & MAINTENANCE

Sow seeds indoors for a jump-start on spring planting.

BACKYARD PROJECTS FOR WINTER

Plan a garden. A successful garden attractive to both humans and wildlife takes time, effort, money and maintenance. Starting with a good plan can help cut down on all of those things.

First, put your ideas on paper. Draw a map of your yard with existing trees, shrubs, slopes, patios and whatever else is established. The map can be formal (a scale version on graph paper) or a casual illustration. Make note of elements that influence the kind of plantings that will follow. Consider factors that will affect how your garden will grow such as sunlight, shade, wind, drainage, access to water, and recreational activity on your property. There are templates and programs online to help with garden design.

Next, choose your plants with guidance from your map and this publication. Budget, ease of care, compatibility with neighboring plants, aesthetics and size all come into play in garden design. A tricky part of native plant gardening is acquiring plants. Research online and local nurseries inquiring about where to obtain the species that interest you. Mail ordering plants and seeds is an option worth considering.

Once you have your map completed and your plants selected, it will soon be spring and time to get your hands dirty! Remember to start small and grow from there! Gardening for butterflies should be looked at as a marathon, not a sprint. Have fun and ease into it! As any experienced green thumb will tell you, there's always room for improvement and new ideas.

COMMON WINTER BLOOMS



COMMON WITCH-HAZEL

Hamamelis virginiana

Yellow, strap-like flowers of this shrub are among the last blooms to appear in the fall. The large, rounded, dark green leaves often hang onto the winter branches. The fruit capsules mature a year after flowering, splitting open to expel seeds. Tolerant of road salt and clay soil and can grow in partial shade. Height 10-15 feet.



COMMON NINEBARK

Physocarpus opulifolius

A cold-hardy, native shrub. Pinkish-white flower clusters in late spring turn into persistent seed capsules and exfoliating bark adds to the season. Papery seed capsules that stay long into the season attract seed-eating birds. This versatile plant can be grown in all soil types with full sun or partial shade. Height 3-10 feet.



SHAGBARK HICKORY

Carya ovata

This open landscape tree is named for its peeling bark giving the tree a shaggy appearance. Hickory leaves are known to support up to 200 different species of caterpillars. Tolerates dry sites and occasional flooding but prefers moist, well-drained soil. Full or partial sun preferred. Height up to 80 feet.



COMMON WINTERBERRY

Ilex verticillata

This easy-growing shrub grows well in wet or dry sites and with full sun or partial shade. The flowers are attractive to butterflies in summer and the berries are attractive to birds in winter. Both male and female plants are required to produce the berries. Height 3- 12 feet.

NECTAR PLANTS



CULVER'S ROOT

Veronicastrum virginicum

Sun: Full

Bloom Period: Mid-Late

Soil: Moist-Mesic

Height: 48-60 inches



JOE PYE WEED

Eutrochium purpureum

Sun: Full

Bloom Period: Mid

Soil: Average

Height: 48-60 inches



GOLDENROD

Solidago spp.

Sun: Full

Bloom Period: Mid-Late

Soil: Average

Height: 30-48 inches



CHOCKECHERRY

Prunus virginiana

Sun: Partial

Bloom Period: Early

Soil: Dry acidic

Height: 20-30 feet



SPIDERWORT

Tradescantia spp.

Sun: Partial

Bloom Period: Early-Mid

Soil: Average

Height: 24-36 inches



PURPLE CONEFLOWER

Echinacea

Sun: Partial

Bloom Period: Mid

Soil: Average

Height: 10-20 inches



INDIGO

Baptisia spp.

Sun: Full

Bloom Period: Early

Soil: Average

Height: 24-60 inches



CARDINAL FLOWER

Lobelia cardinalis

Sun: Partial

Bloom Period: Mid-Late

Soil: Wet

Height: 30-36 inches



CUP PLANT

Silphium perfoliatum

Sun: Full

Bloom Period: Mid-Late

Soil: Moist

Height: 72-96 inches



LUPINE

Lupinus spp.

Sun: Full
Soil: Dry acidic

Bloom Period: Early
Height: 8-24 inches



IRONWEED

Vernonia spp.

Sun: Full
Soil: Average

Bloom Period: Mid-Late
Height: 48-60 inches



LANCELEAF COREOPSIS

Coreopsis lanceolata

Sun: Partial
Soil: Average

Bloom Period: Mid
Height: 24-48 inches



PHLOX

Phlox spp.

Sun: Partial
Soil: Average

Bloom Period: Early
Height: 10-20 inches



BEEBALMS/BERGAMOT

Monarda spp.

Sun: Partial
Soil: Average

Bloom Period: Mid
Height: 30-48 inches



COMMON BONESET

Eupatorium perfoliatum

Sun: Full
Soil: Average

Bloom Period: Mid
Height: 48-60 inches



RATTLESNAKE MASTER

Eryngium yuccifolium

Sun: Partial
Soil: Dry

Bloom Period: Mid-Late
Height: 24-36 inches



OBEDIENT PLANT

Physostegia virginiana

Sun: Partial
Soil: Moist

Bloom Period: Mid-Late
Height: 30-45 inches



BLAZING STARS

Liatris spp.

Sun: Full
Soil: Dry

Bloom Period: Mid-Late
Height: 18-48 inches

EASTERN TIGER SWALLOWTAIL

PAPILIO GLAUCUS



Photo by NINA HARFMANN

The Eastern tiger swallowtail is the most common swallowtail butterfly in Ohio. They love to siphon mineral salts from mud puddles forming “puddle clubs.”

Like most swallowtails they are very strong fliers.

HOST PLANTS: a variety of woody plants, including sassafras (*Sassafras albidum*), spicebush (*Lindera benzoin*), various ash (*Fraxinus* spp.), tulip tree (*Liriodendron tulipifera*), and black cherry (*Prunus serotina*). Trees and shrubs are important species, but keep in mind that they take up more space than typical garden plants.

WINGSPAN: 3.5 inches to 5.5 inches



HOST PLANT PROFILE



SASSAFRAS *Sassafras albidum*

SUN: Full to partial

SOIL: Average

BLOOM PERIOD: Mid

HEIGHT: 30-60 feet

SPICEBUSH SWALLOWTAIL

PAPILIO TROILUS



Photo by FRODE JACOBSEN

Spicebush swallowtails reach peak abundance in large forests of southern Ohio, where it can be numerous. There are normally two broods, in spring and late summer. Because hatches of each brood extend over several weeks, this species can be found from early spring into late fall. The caterpillar is an exceptional example of deceptive camouflage. It appears to have a fearsome snake-like face, which presumably can frighten off potential predators.

HOST PLANTS: spicebush (*Lindera benzoin*), sassafras (*Sassafras albidum*), and members of the laurel family (*Lauraceae*).

WINGSPAN: 3.5 inches to 5.5 inches



HOST PLANT PROFILE



SPICEBUSH *Lindera benzoin*

SUN: Full to partial

SOIL: Average

BLOOM PERIOD: Mid

HEIGHT: 6-12 feet

GIANT SWALLOWTAIL

PAPILIO CRESPHONTES




Photo by KEVIN COLLISON

Numbers are cyclical from year to year, but even in boom years this is Ohio's least common swallowtail. They are usually observed in close proximity to host plant populations. Giant swallowtails are one of the largest butterflies in North America. They are swift, powerful flyers and can quickly vanish to the tree canopy if disturbed.

HOST PLANTS: members of the citrus family (*Rutaceae*). There are only two species in Ohio, prickly-ash (*Zanthoxylum americanum*) and wafer-ash (*Ptelea trifoliata*). Neither species is widely abundant and they tend to form localized colonies. Both plants are more frequent in western Ohio and so is this swallowtail.

WINGSPAN: 4.5 inches to 5.5 inches



HOST PLANT PROFILE



Photo by MARINO DENISENKO

PRICKLY-ASH *Zanthoxylum americanum*

SUN: Full to partial

SOIL: Dry

BLOOM PERIOD: Mid

HEIGHT: 15-20 feet

BLACK SWALLOWTAIL

PAPILIO POLYXENES




Photo by PAUL REEVES

The larvae of black swallowtails are attracted to the oils of plants such as dill, parsley, celery, and carrots. These plants produce specific chemicals that repel insects that try to eat them. Black swallowtail larvae are resistant to these chemicals, so when they feed on the plants, the chemicals make them bad-tasting to bird predators.

HOST PLANTS: native and non-native species in the parsley family (*Apiaceae*). Ohio host plants include Honewort (*Cryptotaenia canadensis*) and garden herbs like fennel (*Foeniculum vulgare*) and dill (*Anethum graveolens*). These non-native species are not considered invasive and will not compete with other garden plants.

WINGSPAN: 2.5 inches to 4.2 inches



HOST PLANT PROFILE



Photo by GERRY BISHOP

GOLD ALEXANDERS *Zizia aurea*

SUN: Full to partial

SOIL: Wet

BLOOM PERIOD: Mid

HEIGHT: 1-2 feet

PEARL CRESCENT

PHYCIODES THAROS




Photo by DAVID BYRON KEENER



Pearl crescents have three to four broods a year, so you can find butterflies on the wing on almost any field outing during the season. They are particularly frequent in old fields with an abundance of asters, and you can occasionally find them gathered in large numbers at a puddle or water source. Like some butterfly species, males are aggressive, darting out to investigate other butterflies and even insects. Pearl crescents overwinter in the larval stage.

HOST PLANTS: many species of Aster, pearl crescents are thought to use the subgenus *Euaster* in particular which includes twenty-two of Ohio's thirty-five aster species.

WINGSPAN: 1.25 inches to 1.6 inches



HOST PLANT PROFILE



Photo by BILDAGENTUR ZONAR GMBH

NEW ENGLAND ASTER *Symphyotrichum novae-angliae*

*Asters are also beneficial nectar plants

SUN: Partial

BLOOM PERIOD: Mid

SOIL: Dry


HEIGHT: 24-60 inches

RED ADMIRAL

VANESSA ATALANTA




Photo by NINA HARFMANN



This species nectars on many flowers and is attracted to mud puddles and animal scat. Most brushfooted butterflies such as the red admiral can be attracted to your backyard by baiting with rotten fruit. It is a fast and erratic flier.

HOST PLANTS: wood nettle (*Laportea canadensis*), stinging nettle (*Urtica dioica*) and Pennsylvania pellitory (*Parietaria pensylvanica*). While nettles aren't a popular choice for gardeners, they are essential for this species. Consider allowing them to grow if they are naturally found on your property.

WINGSPAN: 1.75 inches to 2.5 inches



HOST PLANT PROFILE



Photo by MANFRED RUCKSZIO

NETTLES *Urtica dioica*

SUN: Full
SOIL: Average

BLOOM PERIOD: Mid
HEIGHT: 18-24 inches

QUESTION MARK

POLYGONIA INTERROGATIONIS



Photo by LEENA ROBINSON




Hibernators such as the question mark, comma, and mourning cloak are some of the longest lived butterflies in Ohio.

They can often be seen on sunny, warm, winter days or early spring days. The fall adults hibernate over winter in hollow logs, and in earthen crevices. The males are attracted to mud puddles, tree sap, rotten fruit, animal scat, and carrion.

HOST PLANTS: Hackberry (*Celtis occidentalis*), American elm (*Ulmus americana*), red elm (*Ulmus rubra*), stinging nettle (*Urtica procera*), and false nettle (*Boehmeria cylindrica*).

WINGSPAN: 2.25 inches to 3 inches



HOST PLANT PROFILE




COMMON HACKBERRY *Celtis occidentalis*

SUN: Full to partial

SOIL: Average

BLOOM PERIOD: Mid

HEIGHT: 40-60 feet



AMERICAN LADY

VANESSA VIRGINIENSIS



The ornate inscriptions on the underwing surfaces of this common butterfly are quite striking, as if an abstract artist used the wings as a canvas.

It was once known as the American beauty. This somewhat wary species is best approached when nectaring at a favored flower. This species is first seen on the wing in the early spring, but is more common in June. There are two to three broods per year.

HOST PLANTS: Plantain-leaved pussytoes (*Antennaria plantaginifolia*), fragrant cudweed (*Gnaphalium obtusifolium*), and burdock (*Arctium* species).

WINGSPAN: 1.75 inches to 2.4 inches




Photo by ELIZABETH SPENCER

HOST PLANT PROFILE



Photo by FRITZ FLOHR REYNOLDS


PUSSYTOES *Antennaria plantaginifolia*

SUN: Full

SOIL: Dry

BLOOM PERIOD: Mid

HEIGHT: 6-12 inches



MONARCH

DANAUS PLEXIPPUS



Monarch butterflies are the longest-lived butterfly found in Ohio, with some adults surviving more than ten months. Most monarchs produced in Ohio make an incredible fall migration to high elevation fir forests in Mexico. After overwintering there, they return north in increments, stopping along the way to lay eggs and thus recolonize the southern states. It is the offspring of this first spring brood, and even second or third broods, that returns to Ohio.

HOST PLANTS: up to 13 species of milkweed found in Ohio including common milkweed (*Asclepias syriaca*), swamp milkweed (*A. incarnata*), and butterfly-weed (*A. tuberosa*).

WINGSPAN: 3.5 inches to 4 inches




Photo by NINA HARFMAN

HOST PLANT PROFILE



Photo by NANCY J. ONDRA

SWAMP MILKWEED *Asclepias incarnata*

*Milkweeds are also beneficial nectar plants

SUN: Full to partial

BLOOM PERIOD: Mid

SOIL: Average to moist

HEIGHT: 48-72 inches

GREAT SPANGLED FRITILLARY

SPEYERIA CYBELE



Photo by KENNETH KEIFER



This is a large butterfly with a strong flight. They frequently stop at flowers, especially milkweeds. Males often gather at mud puddles and animal scat, taking in mineral salts and proteins. Females are larger and darker than males, and do not appear until later in the summer when they lay eggs near host plants – but often not on the plants. Newly hatched larvae overwinter without feeding. In spring, they must make their way to the hosts, upon which they only feed at night.

HOST PLANTS: violets (*Viola spp.*), likely many of the twenty-six native species known to Ohio.

WINGSPAN: 2.9 inches to 3.8 inches



HOST PLANT PROFILE



Photo by MELINDA FAWVER

COMMON BLUE VIOLET *Viola sororia*

SUN: Full to partial

BLOOM PERIOD: Mid

SOIL: Average to moist

HEIGHT: 4-6 inches



ENODIA ANTHEDON

Photo by TIM DANIEL

NORTHERN PEARLY EYE

Northern pearly-eyes get their name from the ringed eye-spots on the ventral hindwing. It is one of the few Ohio butterflies that strictly inhabits deep, shady wooded habitats. Their dark coloration and striped and spotted wing pattern allow them to blend well with their surroundings. When disturbed, they shoot off in an erratic flight that is hard to follow among the trees and dappled sunlight of the forest. This species rarely visits flowers as they typically visit tree sap, rotting fruit, and animal remains.

A number of species of grasses, especially bottlebrush grass (*Elymus hystrix*), river oats (*Chasmanthium latifolium*), longawned wood grass (*Brachyelytrum erectum*), and white grass (*Leersia virginica*).



HOST PLANT PROFILE

BOTTLEBRUSH GRASS


Elymus hystrix

Sun: Full to partial

Soil: Average to dry

Bloom Period: Late

Height: 24-36 inches



EPARGYREUS CLARUS

SILVER-SPOTTED SKIPPER

If only all of our skippers were so easy to identify! Big, bold, and distinctively marked, the common and wide-ranging silver-spotted skipper is far more conspicuous than most of the other skipper species found in Ohio. It, like almost all of Ohio's skippers, has a curved tip to its antennae called an apiculus. Silver-spotteds have a strong and fast flight pattern. They can be hard to follow, but often stop to take nectar at flowers of milkweeds, thistles and other blooming plants.

HOST PLANTS: various species in the pea family (*Fabaceae*), including partridge pea (*Chamaecrista fasciculata*), black locus (*Robinia pseudoacacia*), and honey-locust (*Gleditsia triacanthos*).



HOST PLANT PROFILE

PARTRIDGE PEA


Chamaecrista fasciculata

Sun: Full

Soil: Average to dry

Bloom Period: Mid

Height: 12-36 inches





**CELASTRINA LADON/
CELASTRINA NEGLECTA**

Photo by JOHANN VILORIA



AGLAIS MILBERTI

Photo by FRODE JACOBSEN

SPRING & SUMMER AZURE

These species are extremely similar and until recently were considered to be different forms of the same species. Both are very common in Ohio and sometimes form huge “puddle clubs” at favored muddy spots. The spring azure is one of the first non-hibernating butterflies to appear each spring. Occasionally, it can be found near the Ohio River as early as mid-March.

HOST PLANTS: black cherry (*Prunus serotina*), flowering dogwood (*Cornus florida*), and a variety of other woody plants. Summer azures use wingstem (*Verbesina alternifolia*).



HOST PLANT PROFILE

BLACK CHERRY


Prunus serotina

Sun: Full to partial

Soil: Average

Bloom Period: Mid

Height: 50-80 feet



27

MILBERT'S TORTOISESHELL

Milbert's tortoiseshell, also known as the fire-rim tortoiseshell, has a wingspan between 1.5 and 2.5 inches with the forewing tips squared off. This is a fast butterfly that flits rapidly around woodland roads. The Milbert's tortoiseshell's range includes all of Canada south of the tundra, western U.S., and most of the east. In these regions they are common in wet areas, including moist pastures, marshes, most trails, and roadsides.

When it lands, it may open its wings, often on trees or rocks. From May to October adults mate and lay eggs. The caterpillars congregate in a small web similar to tent caterpillars (moth), but do not leave the web to feed.



HOST PLANT PROFILE

STINGING NETTLE


Urtica dioica

Sun: Full to partial

Soil: Average-moist

Bloom Period: Mid

Height: 3-7 feet





PUBLICATION FUNDING

Funding for this publication was provided by donations to the state income tax checkoff program, sales of the cardinal license plate, and Ohio Wildlife Legacy Stamp.

See below for ways to support free publications like this:

PURCHASE A LICENSE PLATE

Visit your local registrar's office or call the BMV at **1-888-PLATES3**

PURCHASE A LEGACY STAMP

Call the Division of Wildlife at **1-800-WILDLIFE** or visit the web at **wildohiostamp.gov**

MAKE A DONATION

Mail a check to:
Wildlife Diversity Fund
2045 Morse Road Bldg G.
Columbus, OH 43229-6693

Donate through your Ohio tax form:
Go to the second page of the **Ohio IT 1040** income tax form for the tax checkoff program.

For more information about Ohio's native wildlife, please contact the Ohio Division of Wildlife:

1-800-WILDLIFE

(1-800-750-0750 Ohio Relay TTY only)

wildohio.gov

GOVERNOR, STATE OF OHIO
MIKE DEWINE

DIRECTOR, OHIO DEPARTMENT
OF NATURAL RESOURCES
MARY C. MERTZ

CHIEF, DIVISION OF WILDLIFE
KENDRA S. WECKER

PUBLICATION 5089 (0120)

Total Quantities Printed: 12,500 Unit cost: \$0.000 Publication date: 01/20

OHIO DEPARTMENT OF NATURAL RESOURCES DIVISION OF WILDLIFE



 **Ohio Pollinator**
Habitat Initiative

MISSION STATEMENT

To conserve and improve fish and wildlife resources and their habitats for sustainable use and appreciation by all.

The ODNR Division of Wildlife is the state agency responsible for managing Ohio's fish and wildlife resources. The primary source of funding for the division comes from the sale of hunting and fishing licenses, federal excise taxes on hunting, fishing, and shooting equipment, and donations from the public. We care about all wildlife and maintaining stable, healthy wildlife populations. Our challenge is to balance the needs of wildlife, habitat, and people.